

A New Formation Mechanism for the Hottest Horizontal-Branch Stars

Allen V. Sweigart

NASA Goddard Space Flight Center, Code 681, Greenbelt, MD 20771

Thomas M. Brown

Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218

Thierry Lanz, Wayne B. Landsman, Ivan Hubeny

NASA Goddard Space Flight Center, Code 681, Greenbelt, MD 20771

Abstract. Stars with very large mass loss on the red-giant branch can undergo the helium flash while descending the white-dwarf cooling curve. Under these conditions the flash convection zone will mix the hydrogen envelope with the hot helium-burning core. Such “flash-mixed” stars will arrive on the extreme horizontal branch (EHB) with helium- and carbon-rich envelopes and will lie at higher temperatures than the hottest canonical (i.e., unmixed) EHB stars. Flash mixing provides a new evolutionary channel for populating the hot end of the EHB and may explain the origin of the high gravity, helium-rich sdO and sdB stars.

1. Evolution of Hot Flashers

Castellani & Castellani (1993) first showed that stars with large mass loss on the red-giant branch (RGB) can evolve to high effective temperatures before igniting helium in their cores. We have studied the properties of these “hot flashers” by following their evolution through the helium flash to the zero-age horizontal branch (ZAHB) for various amounts of mass loss (Brown et al. 2001; Sweigart et al. 2002). Two examples of our results are given in Figure 1. The “early hot flasher” in the left panel ignites helium between the tip of the RGB and the top of the white dwarf (WD) cooling curve. Mixing between the hydrogen envelope and helium core does not occur in such stars due to the high entropy of the hydrogen shell. Early hot flashers will thus have canonical, hydrogen-rich envelopes on the EHB.

The evolution is dramatically different for the “late hot flasher” in the right panel of Figure 1. Due to the weaker hydrogen shell the flash convection zone will then penetrate deeply into the stellar envelope. Protons from the envelope will be mixed into the helium core and rapidly burned, while helium and carbon from the core will be mixed outward to the stellar surface. Such flash mixing is a consequence of the basic properties of the stellar models and should therefore occur whenever a star ignites helium on the WD cooling curve.




Figure 1. Evolution of an early hot flasher (left panel) and a late hot flasher (right panel) from the main sequence (MS) through the helium flash to the ZAHB. The flash peak is marked (*). Flash mixing began at the end of the late hot flasher track (+).

Flash mixing has many implications for the properties of the EHB stars:

- Flash mixing produces hot EHB stars with greatly enhanced envelope helium and carbon abundances.
- Flash-mixed EHB stars will appear subluminescent in the UV due to their redistributed far-UV flux and larger bolometric corrections, in agreement with the subluminescent EHB stars observed in ω Cen and NGC 2808.
- Flash mixing leads to a sharp dichotomy in the properties of the EHB stars which may explain the gap within the EHB found in optical color-magnitude diagrams of NGC 2808.
- Flash mixing on the WD cooling curve provides a new evolutionary channel for producing hydrogen-deficient stars, particularly the high gravity, helium-rich sdO stars and the minority of sdB stars that are helium-rich.
- The high carbon abundance in the envelopes of the flash-mixed stars may be important for driving the pulsations of the variable sdB stars.

References

- Brown, T. M., Sweigart, A. V., Lanz, T., Landsman, W. B., & Hubeny, I. 2001, *ApJ*, 562, 368
- Castellani, M., & Castellani, V. 1993, *ApJ*, 407, 649
- Sweigart, A. V., Brown, T. M., Lanz, T., Landsman, W. B., & Hubeny, I. 2002, in *ω Centauri: A Unique Window into Astrophysics*, ed. F. van Leeuwen, J. D. Hughes, & G. Piotto (San Francisco: ASP), 261